

REVIEW ARTICLE

ENVIRONMENTAL AGGRESSORS AND SKIN AGING: UNRAVELING THE IMPACT OF EXTERNAL FACTORS ON PHOTODAMAGE

Deepak Joshi^{1*}, Pawandeep Shukla², Naveen K Choudhary³

¹Shri Vaishnav Vidyapeeth Vishwavidyalaya (SVVV), Indore, India

²Shri Vaishnav Vidyapeeth Vishwavidyalaya (SVVV), Indore, India

³B R Nahata College of Pharmacy, Mandsaur, India

Corresponding Author: Deepak Joshi, ORCID Id: 0009-0005-4108-7098

Email Id: djoshi25694@gmail.com


Received on: 02/07/2025, Accepted on: 10/10/2025, Published on: 29/12/2025

ABSTRACT

Skin aging is a complex biological process influenced by both intrinsic and extrinsic factors. While intrinsic aging is determined by genetic and chronological changes, extrinsic aging is accelerated by environmental factors, leading to premature skin damage. Among these, ultraviolet (UV) radiation, air pollution, climate variations, and lifestyle habits such as smoking and poor nutrition contribute significantly to photodamage and skin deterioration. These external aggressors trigger oxidative stress, inflammation, collagen degradation, and DNA damage, ultimately resulting in wrinkles, loss of elasticity, and pigmentation disorders.

This review explores the impact of key environmental factors on skin aging, emphasizing their mechanisms and long-term effects. Additionally, it highlights various preventive strategies, including the use of sunscreens, antioxidants, and lifestyle modifications to mitigate photodamage. By understanding these influences, effective interventions can be developed to maintain skin health and delay the aging process.

Keyword: Skin aging, Environmental factors, UV radiation, Air pollution, Oxidative stress



Graphical Abstract: Impact of External Factors on Photodamage

1. INTRODUCTION

Skin aging is a natural and gradual biological process that affects the structure and function of the skin over time. It is characterized by visible changes such as wrinkles, fine lines, loss of elasticity, uneven pigmentation, and dryness. This process occurs due to both intrinsic and extrinsic factors, with the latter playing a significant role in premature aging.[1]

Photodamage, often referred to as photoaging, specifically results from chronic exposure to ultraviolet (UV) radiation. Unlike chronological aging, which is an inevitable consequence of time, photodamage leads to accelerated

deterioration of the skin's appearance and function. The primary manifestations of photodamage include deep wrinkles, hyperpigmentation, reduced skin elasticity, and rough texture. Over time, excessive UV exposure also increases the risk of skin malignancies, making it a serious dermatological concern. [2-3]

Skin aging is broadly categorized into intrinsic aging and extrinsic aging, both of which contribute to the overall aging process but differ in their underlying causes and mechanisms. [4]

Intrinsic aging, also known as chronological aging, is a genetically programmed process influenced by the

body's natural metabolic activities. It results from physiological changes that occur over time, including reduced collagen and elastin production, slower cell turnover, and diminished sebaceous gland activity. The skin gradually loses its ability to retain moisture, leading to dryness and fine wrinkles. Additionally, the decrease in fibroblast activity weakens the skin's structural integrity, causing sagging and thinning over time. Unlike extrinsic aging, intrinsic aging is inevitable and progresses at a genetically predetermined rate. [5]

Extrinsic aging, in contrast, is caused by external environmental factors that accelerate the aging process beyond what is dictated by genetics. Major contributors to extrinsic aging include:

- **Ultraviolet (UV) Radiation:** Chronic sun exposure damages skin cells through oxidative stress, DNA mutations, and collagen breakdown, leading to photodamage.
- **Air Pollution:** Pollutants such as particulate matter, heavy metals, and toxins contribute to inflammation, free radical production, and disruption of the skin barrier.
- **Smoking:** Tobacco smoke generates harmful free radicals that degrade collagen and elastin, leading to premature wrinkles and dull skin.
- **Climate Variations:** Extreme temperatures, humidity fluctuations, and harsh weather conditions impact skin hydration and barrier function.

- **Diet and Lifestyle Factors:** Poor nutrition, excessive alcohol consumption, and lack of sleep can impair the skin's natural repair mechanisms and exacerbate aging.

Among these factors, UV radiation is considered the most potent extrinsic factor responsible for skin aging. Other elements, such as pollution and lifestyle choices, further compound the effects of UV exposure, leading to accelerated deterioration of skin health. [6]

The study of environmental influences on skin aging is crucial due to the rising global exposure to pollutants, climate changes, and increased UV radiation resulting from ozone layer depletion. Over the years, dermatological research has emphasized the role of oxidative stress, inflammation, and cellular damage in extrinsic aging. Understanding these mechanisms is essential for developing effective preventive strategies such as advanced skincare formulations, antioxidant-rich diets, and public awareness campaigns on photoprotection. [7]

Additionally, the increasing demand for anti-aging interventions has led to the development of targeted dermatological solutions, including broad-spectrum sunscreens, topical antioxidants (such as vitamins C and E), and novel bioactive compounds aimed at repairing and protecting the skin from environmental damage. By studying these external

influences, researchers and healthcare professionals can help individuals adopt better skincare practices, thereby reducing premature aging and improving overall skin health. [8]

2. ENVIRONMENTAL FACTORS CONTRIBUTING TO SKIN AGING AND PHOTODAMAGE

Skin aging is significantly influenced by environmental factors that accelerate damage to the skin's structure and function. These external aggressors induce oxidative stress, inflammation, and degradation of collagen and elastin, leading to premature aging and photodamage. The most prominent environmental factors contributing to skin aging include ultraviolet (UV) radiation, air pollution, smoking, climate conditions, and dietary habits. [9-11]

2.1. Ultraviolet (UV) Radiation

Ultraviolet (UV) radiation is one of the most significant environmental factors contributing to extrinsic skin aging and photodamage. Prolonged exposure to UV rays accelerates the aging process through mechanisms such as oxidative stress, DNA damage, collagen degradation, and elastin fiber breakdown. These effects collectively lead to wrinkles, pigmentation disorders, loss of skin elasticity, and an increased risk of skin cancer. [12-15]

2.1.1 UV-Induced Oxidative Stress and DNA Damage

One of the primary ways UV radiations contributes to skin aging is by generating reactive oxygen species (ROS), which induce oxidative stress at the cellular level. These ROS attack lipids, proteins, and DNA, leading to cellular dysfunction and structural deterioration.

Oxidative Stress Mechanism:

- UV exposure increases ROS production, depleting the skin's natural antioxidants such as vitamin C, vitamin E, and glutathione.
- Excessive ROS cause lipid peroxidation, damaging cell membranes and leading to loss of moisture and barrier dysfunction.
- ROS activate matrix metalloproteinases (MMPs), enzymes that degrade collagen and elastin fibers, weakening the skin's structural integrity.

DNA Damage and Mutations:

- UVB radiation directly damages nuclear and mitochondrial DNA, forming cyclobutane pyrimidine dimers (CPDs) and 6-4 photoproducts, leading to mutations and cellular apoptosis.
- Accumulation of damaged DNA can lead to cellular senescence, where cells lose their ability to divide and repair, accelerating the aging process.

- In severe cases, DNA mutations increase the risk of skin cancers, including basal cell carcinoma, squamous cell carcinoma, and melanoma.

2.1.2 Role of UVA vs. UVB in Premature Aging (Photoaging)

UV radiation is divided into three types: UVA (320–400 nm), UVB (280–320 nm), and UVC (100–280 nm). UVC is mostly absorbed by the ozone layer, while UVA and UVB are the primary contributors to skin aging and photodamage. [16-19]

Table 1: Comparison of UVA and UVB in Skin Aging

Factor	UVA	UVB
Penetration	Deep (dermis)	Superficial (epidermis)
DNA Damage	Indirect via oxidative stress	Direct via pyrimidine dimer formation
Effects on Skin	Collagen and elastin breakdown, long-term aging	Sunburn, hyperpigmentation, and direct mutation risk
Cancer Risk	Contributes to melanoma and non-melanoma cancers	High risk for basal and squamous cell carcinoma
Daily Exposure Risk	Present throughout the day, even through glass	Strongest midday, less through windows

2.1.3 Effects on Collagen Degradation and Elastin Fibers

Collagen and elastin are the key structural proteins responsible for maintaining skin firmness, elasticity, and resilience. Chronic UV exposure accelerates their degradation, leading to wrinkling, loss of firmness, and sagging skin. [20]

Collagen Degradation

Collagen is the primary structural protein in the dermis, responsible for maintaining skin strength and smoothness. UV

radiation reduces collagen synthesis while increasing collagen degradation through the following mechanisms:

- Activation of Matrix Metalloproteinases (MMPs):

- MMP-1 (collagenase) and MMP-3 break down collagen types I and III, essential for skin firmness.
- UVA exposure upregulates MMP activity, leading to rapid breakdown of collagen fibers.

- Reduction in Collagen Production:

- UV radiation suppresses fibroblast activity, the key cells responsible for collagen synthesis.
- Leads to a gradual thinning of the dermis, making the skin more fragile and prone to wrinkles.

Elastin Fiber Damage

Elastin fibers allow the skin to stretch and recoil, giving it flexibility. Chronic UV exposure leads to solar elastosis, a condition characterized by thickened, damaged elastin fibers that lose their normal function. [21]

- Solar Elastosis Features:

- Abnormal accumulation of disorganized elastin fibers in the dermis.
- Causes skin roughness, leathery texture, and loss of elasticity.
- Appears as deep wrinkles, crepey skin, and sagging in sun-exposed areas like the face, neck, and hands.

2.2. Air Pollution

Air pollution significantly impacts skin health through various pollutants, including particulate matter (PM), polycyclic aromatic hydrocarbons (PAHs), and heavy metals. These pollutants can penetrate the skin, leading to oxidative stress, inflammation, and disruption of the skin barrier function[22-24]

2.2.1 Pollutants and Their Effects

- **Particulate Matter (PM):** Particularly PM_{2.5}, is linked to skin diseases such as atopic dermatitis and psoriasis, causing oxidative stress and inflammatory responses.
- **Polycyclic Aromatic Hydrocarbons (PAHs):** These compounds exacerbate skin conditions by inducing inflammation and oxidative damage, contributing to premature aging and skin cancers.
- **Heavy Metals:** Exposure to heavy metals can lead to skin barrier dysfunction and inflammatory skin diseases, further complicating skin health.

2.2.2 Mechanisms of Damage

Oxidative Stress: Reactive oxygen species (ROS) produced by pollutants harm cellular constituents and activate inflammatory pathways like MAPK and NF- κ B. **Inflammation:** Exposure to pollutants causes an increase in the production of inflammatory cytokines, which exacerbates diseases including dermatitis and acne. **Skin Barrier Function:** When pollutants interfere with the skin's defenses, it becomes more permeable and vulnerable to environmental stimuli. [25-28]

2.3. Climate Conditions (Temperature & Humidity)

Extreme temperatures and humidity significantly impact skin structure and function, particularly affecting the skin barrier and hydration levels. Cold and low humidity conditions lead to decreased skin barrier function, increased trans epidermal water loss (TEWL), and heightened susceptibility to irritants and allergens, exacerbating conditions like atopic dermatitis. Conversely, high humidity can disrupt permeability homeostasis, reducing the synthesis of lipids and natural moisturizing factors, which are crucial for maintaining skin hydration and barrier integrity. Proper hydration is essential for skin health, as it regulates temperature and facilitates tactile interaction with the environment; thus, maintaining an optimal balance of humidity is vital for preserving skin barrier function. Overall, both extreme temperatures and humidity levels play critical roles in skin health, necessitating appropriate environmental management to mitigate adverse effects. [29-32]

2.4. Smoking and Tobacco Exposure

Smoking significantly impacts skin health through mechanisms involving free radical generation, collagen breakdown, and alterations in skin elasticity and pigmentation. Tobacco smoke induces oxidative stress, leading to increased production of reactive oxygen species (ROS) that damage skin cells and

extracellular matrix components, particularly collagen and elastin, which are crucial for maintaining skin structure and elasticity. This oxidative damage is associated with a decrease in skin elasticity, as evidenced by studies linking reduced skin viscoelasticity to pulmonary emphysema and inflammation in smokers. Furthermore, smoking contributes to irregular pigmentation and accelerated skin aging, characterized by wrinkles and a loss of hydration due to impaired blood flow and nutrient delivery. Although some studies suggest that smoking does not significantly alter the morphology of elastic fibers, the overall detrimental effects on skin quality and appearance are well-documented. Thus, smoking exacerbates skin aging through complex biochemical pathways that compromise skin integrity and aesthetics[33]

2.5. Diet and Nutrition

Diet and nutrition significantly influence skin health, particularly through the roles of antioxidants, vitamins, and hydration, as well as the impact of processed foods and sugar on skin glycation. Antioxidants and vitamins, found in fruits, vegetables, and nuts, are essential for maintaining skin integrity and combating oxidative stress, which is linked to aging and skin damage. Adequate hydration is also crucial, as it supports skin elasticity and overall health. Conversely, excessive sugar intake and processed foods contribute to skin glycation, leading to the formation of advanced glycation end-products (AGEs)

that damage collagen and elastin, exacerbating skin aging and inflammatory conditions. A low-glycemic index diet can mitigate these effects, promoting healthier skin by reducing glycation and inflammation. Thus, adopting a balanced diet rich in nutrients while minimizing processed foods is vital for optimal skin health. [34]

2.6. Other Environmental Factors

Blue light exposure from screens and artificial lighting has been increasingly scrutinized for its potential impact on skin health and aging. While the sun remains the primary source of blue light, artificial sources such as LEDs and electronic devices contribute significantly to cumulative exposure, which can lead to oxidative stress and skin damage, particularly in individuals with hypersensitivity. This exposure is also implicated in age-related macular degeneration, as indoor lighting and screen exposure, although less intense than sunlight, can still pose risks due to prolonged exposure times. Infrared radiation, another environmental factor, contributes to skin aging by promoting oxidative stress and inflammation, which accelerates the degradation of skin structure and function. Lifestyle factors, including stress and sleep deprivation, further exacerbate aging processes. Stress is linked to increased oxidative stress and inflammation, which can accelerate telomere attrition and aging-related disorders. Sleep deprivation similarly

affects skin health by impairing repair mechanisms and increasing susceptibility to environmental stressors. Collectively, these factors underscore the complex interplay between environmental exposures and lifestyle choices in the aging process, highlighting the need for comprehensive strategies to mitigate their effects. [35-39]

3. MECHANISMS OF ENVIRONMENTAL SKIN DAMAGE

Environmental skin damage is primarily driven by mechanisms such as oxidative stress, inflammation, and matrix metalloproteinase (MMP) activation, leading to collagen degradation and DNA damage. Ultraviolet (UV) radiation, particularly UVA and UVB, generates reactive oxygen species (ROS) that induce oxidative stress, resulting in DNA lesions and inflammatory responses that can promote skin aging and cancer. MMPs, activated by oxidative stress, contribute to collagen breakdown, further compromising skin integrity. Additionally, environmental pollutants exacerbate these effects by disrupting the skin barrier and increasing ROS production, which is linked to various skin disorders, including inflammatory diseases and carcinogenesis. The interplay between oxidative stress, inflammation, and DNA damage underscores the complexity of environmental impacts on skin health, necessitating targeted therapeutic strategies to mitigate these effects. [40-41]

4. PREVENTION AND PROTECTIVE STRATEGIES

4.1. Sunscreens and Photoprotection

Sunscreens and photoprotection are essential strategies for mitigating the harmful effects of ultraviolet (UV) radiation, which is a significant contributor to photoaging and photo carcinogenesis. Effective photoprotection encompasses a range of methods, including sun avoidance, protective clothing, and the application of broad-spectrum sunscreens that shield against both UVA and UVB rays. Regular use of these sunscreens is crucial, as they not only prevent immediate skin damage, such as sunburn, but also protect cellular DNA from mutations that can lead to skin cancer. Emerging technologies in sunscreen formulation are enhancing their efficacy, with new filters providing broader protection, including against visible light, which has been linked to conditions like hyperpigmentation. Overall, a comprehensive approach combining sunscreen use with other protective measures is vital for maintaining skin health and preventing premature aging. [42-45]

4.2. Antioxidants and Skincare Interventions

Antioxidants play a crucial role in skincare interventions aimed at preventing oxidative stress-related skin damage, including photoaging and skin cancer. Key antioxidants such as Vitamins C and E, retinol, and polyphenols from green tea

have demonstrated significant protective effects against UV-induced oxidative stress, which is a primary contributor to skin aging and carcinogenesis. These compounds work synergistically to enhance collagen synthesis, reduce inflammation, and promote skin cell regeneration. Additionally, dietary antioxidants have shown chemo-preventive effects against skin cancer and efficacy in treating various dermatological conditions, including rosacea and acne. Comprehensive skincare strategies that combine topical applications with dietary intake of antioxidants, alongside preventive measures like sun protection, are essential for maintaining skin health and mitigating age-related changes. [46-47]

4.3. Lifestyle Modifications

Lifestyle modifications constitute an essential component in the prevention, management, and even reversal of many chronic conditions, including skin-related disorders and premature skin aging. These modifications extend beyond simple recommendations and require a comprehensive approach tailored to each individual's health status and risk factors.

A balanced diet is fundamental, particularly one abundant in fruits, vegetables, whole grains, lean proteins, and healthy fats, as these provide vitamins, minerals, and antioxidants that protect the skin from oxidative stress. Adequate hydration further supports epidermal

function and improves skin texture. Limiting processed foods, sugar, and excessive alcohol consumption helps reduce systemic inflammation, which can otherwise accelerate dermatological issues.

Physical activity serves multiple benefits not only in improving cardiovascular health but also in enhancing skin physiology. Regular exercise promotes blood circulation, oxygen delivery, and nutrient supply to skin tissues, improving firmness, elasticity, and wound-healing capacity. Moderate exercise is also associated with reduced systemic inflammation, benefiting both skin quality and overall vitality.

Stress management is another cornerstone of lifestyle modification. Chronic psychological stress has been linked to inflammatory skin disorders such as psoriasis, eczema, and acne. Incorporating relaxation techniques—such as yoga, meditation, deep-breathing exercises, or mindfulness practices—can correct hormonal imbalances and attenuate stress-related skin flare-ups.

Sleep hygiene plays a critical role in skin recovery, as restorative sleep regulates hormonal cycles, promotes collagen synthesis, and facilitates cellular repair. Chronic sleep deprivation not only contributes to dullness and early wrinkling of the skin but also weakens the immune response, increasing vulnerability to infections and inflammatory conditions. Establishing consistent sleep schedules and

limiting screen exposure before bedtime significantly support skin regeneration.

Lifestyle changes must also include avoiding harmful habits and adopting protective measures. Smoking cessation is essential since tobacco accelerates skin aging through collagen degradation and reduced blood flow. Similarly, excessive sun exposure remains one of the most significant extrinsic aging factors; hence, the use of broad-spectrum sunscreens, protective clothing, and avoidance of tanning practices are indispensable.

It is important to note that the efficacy of lifestyle modifications often depends on individual adherence and personalization. Factors such as age, genetics, cultural practices, and socioeconomic background influence these adjustments. Therefore, structured healthcare guidance, behavioral support, and long-term follow-up are often necessary to ensure effectiveness and sustainability of these interventions. [48-50]

5. FUTURE PERSPECTIVES AND CHALLENGES

The future of dermatological science, particularly in the realm of anti-aging and skincare, is marked by rapid technological advancements and growing personalization. Ongoing research into cellular mechanisms such as oxidative stress, DNA damage, and cellular senescence is leading to novel anti-aging therapies, including the use of stem cells, growth factors, and gene-editing

technologies like CRISPR-Cas9. Alongside these biological innovations, personalized skincare based on environmental exposure is gaining momentum. By leveraging wearable sensors, exposome data, and AI-driven analytics, skincare regimens can now be tailored to individual needs influenced by external factors such as UV radiation, pollution, and climate. Furthermore, the incorporation of biotechnology and nanotechnology is transforming transdermal drug delivery, with nanosystems like liposomes, dendrimers, and invasomes enhancing skin penetration and therapeutic efficacy. Bioengineered skin models and 3D bioprinting also hold promise for product testing and skin regeneration. Despite these exciting developments, challenges such as regulatory approval, safety concerns, cost, and scalability must be addressed to ensure these innovations transition successfully from the lab to clinical and consumer applications. [51-53]

6. CONCLUSION

This study provides new insights into the multifaceted nature of skin aging, emphasizing both intrinsic biological processes and the wide-ranging impact of the external environment. By systematically examining how environmental exposures such as UV radiation, pollution, and climate variability accelerate skin aging, the research underscores the urgency of comprehensive preventive strategies. The novel integration of emerging concepts—such as the

exposome, epigenetic markers, and the microbiome—demonstrates the growing importance of personalized and adaptive care in dermatology. These findings pave the way for the future development of innovative diagnostic tools and individualized interventions, thereby holding promise for more effective, targeted management of skin aging. This work not only enriches current scientific understanding but also offers a foundation for future research in precision dermatology and preventive skin health.

FUNDING: This study did not receive any funding from an agency or person.

REFERENCES

- [1] McDaniel, D., Farris, P., & Valacchi, G. (2018). Atmospheric skin aging Contributors and inhibitors. *Journal of Cosmetic Dermatology*, 17(1), 124–137. <https://doi.org/10.1111/jocd.12418>
- [2] Pilkington, S. M., Bulfone-Paus, S., Griffiths, C. E. M., & Watson, R. E. B. (2021). Inflammaging and the skin. *Journal of Investigative Dermatology*, 141(5), 1087–1095. <https://doi.org/10.1016/j.jid.2020.10.039>
- [3] Pecorelli, A., Woodby, B., Prieux, R., Valacchi, G., & Nrf2. (2019). Involvement of 4-hydroxy-2-nonenal in pollution-induced skin damage. *BioFactors*, 45(4), 536–547. <https://doi.org/10.1002/biof.1510>

- [4] Ferrara, F., Prioux, R., Woodby, B., Pecorelli, A., & Valacchi, G. (2021). Inflammasome activation in pollution-induced skin conditions. *Plastic and Reconstructive Surgery*, 147(1S), 15S–24S. <https://doi.org/10.1097/PRS.00000000000007557>
- [5] Martic, I., Jansen-Dürr, P., & Cavinato, M. (2022). Effects of air pollution on cellular senescence and skin aging. *Cells*, 11(14), 2220. <https://doi.org/10.3390/cells11142220>
- [6] Qiao, L., Cai, J., Wang, H., Wang, L., Chen, R., Kan, H., & Zhang, Y. (2014). PM_{2.5} constituents and hospital emergency-room visits in Shanghai, China. *Environmental Science & Technology*, 48(19), 10406–10414. <https://doi.org/10.1021/es502119b>
- [7] Kousha, T., & Valacchi, G. (2015). The air quality health index and emergency department visits for urticaria in Windsor, Canada. *Journal of Toxicology and Environmental Health, Part A*, 78(8), 524–533. <https://doi.org/10.1080/15287394.2015.1007146>
- [8] Lyngsie, G., Krumina, L., Tunlid, A., & Persson, P. (2018). Generation of hydroxyl radicals from reactions between a dimethoxyhydroquinone and iron oxide nanoparticles. *Scientific Reports*, 8, 10834. <https://doi.org/10.1038/s41598-018-29204-4>
- [9] Ferreira, M. S., Magalhães, M. C., Oliveira, R., & Guimarães, M. (2021). Trends in the use of botanicals in anti-aging cosmetics. *Molecules*, 26(12), 3584. <https://doi.org/10.3390/molecules26123584>
- [10] Pambianchi, E., Ferrara, F., Pecorelli, A., Woodby, B., & Valacchi, G. (2020). Blueberry extracts as a novel approach to prevent ozone-induced cutaneous inflammasome activation. *Oxidative Medicine and Cellular Longevity*, 2020, 9571490. <https://doi.org/10.1155/2020/9571490>
- [11] Saeedi, M., Eslamifar, M., & Khezri, K. (2019). Kojic acid applications in cosmetic and pharmaceutical preparations. *Biomedicine & Pharmacotherapy*, 110, 582–593. <https://doi.org/10.1016/j.biopha.2018.11.100>
- [12] Rathee, P., Kumar, S., Kumar, D., & Arya, R. (2021). Skin hyperpigmentation and its treatment with herbs: An alternative method. *Future Journal of Pharmaceutical Sciences*, 7(1), 132. <https://doi.org/10.1186/s43094-021-00258-5>
- [13] Jin, S. P., Li, Z., Choi, E. K., Lee, S., Kim, Y. K., Seo, E. Y., & Chung, J. H. (2018). Urban particulate matter in air pollution penetrates into the barrier-disrupted skin and produces ROS-dependent cutaneous inflammatory response in vivo. *Journal of Dermatological Science*, 91(2),

- 175–183. <https://doi.org/10.1016/j.jdermsci.2018.04.014>
- [14] Mancebo, S. E., & Wang, S. Q. (2015). Recognizing the impact of ambient air pollution on skin health. *Journal of the European Academy of Dermatology and Venereology*, 29(12), 2326–2332. <https://doi.org/10.1111/jdv.13252>
- [15] Soeur, J., Belaïdi, J. P., Chollet, C., et al. (2017). Photo-pollution stress in skin: Traces of pollutants (PAH and particulate matter) impair redox homeostasis in keratinocytes exposed to UVA1. *Journal of Dermatological Science*, 86(2), 162–169. <https://doi.org/10.1016/j.jdermsci.2017.02.002>
- [16] Latha, M. S., Martis, J., Shobha, V., et al. (2013). Sunscreening agents: A review. *Journal of Clinical and Aesthetic Dermatology*, 6(1), 16–26.
- [17] Heurung, A. R., Raju, S. I., & Warshaw, E. M. (2014). Adverse reactions to sunscreen agents: Epidemiology, responsible irritants and allergens, clinical characteristics, and management. *Dermatitis*, 25(6), 289–326. <https://doi.org/10.1097/DER.0000000000000061>
- [18] Ruszkiewicz, J. A., Pinkas, A., Ferrer, B., Peres, T. V., Tsatsakis, A., & Aschner, M. (2017). Neurotoxic effect of active ingredients in sunscreen products: A contemporary review. *Toxicology Reports*, 4, 245–259. <https://doi.org/10.1016/j.toxrep.2017.05.007>
- [19] Kim, S., & Choi, K. (2014). Occurrences, toxicities, and ecological risks of benzophenone-3, a common component of organic sunscreen products: A mini-review. *Environment International*, 70, 143–157. <https://doi.org/10.1016/j.envint.2014.05.015>
- [20] Shah, K., & Murray, P. G. (Inventors). (2015, July 27). Process for producing a coated powder (European Patent No. 3237548).
- [21] Herrling, T., Jung, K., & Fuchs, J. (2006). Measurements of UV-generated free radicals/reactive oxygen species (ROS) in skin. *Spectrochimica Acta Part A: Molecular and Biomolecular Spectroscopy*, 63(4), 840–845. <https://doi.org/10.1016/j.saa.2005.09.046>
- [22] Sarkas, H. W., Cureton, K., & Jung, K. (2018). The photophysics of environmental protection for preventing premature aging. *Euro Cosmetics*, 26, 20–25.
- [23] Halliday, G. M., Damian, D. L., Rana, S., & Byrne, S. N. (2012). The suppressive effects of ultraviolet radiation on immunity in the skin and internal organs: Implications for autoimmunity. *Journal of Dermatological Science*, 66(3),

- 176–182. <https://doi.org/10.1016/j.jdermsci.2012.05.001>
- [24] NS Phytomoist. (2019, May 16). Next Step Laboratories. <https://www.nextsteplab.com/ns-phytomoist>
- [25] Puri, P., Nandar, S. K., Kathuria, S., & Ramesh, I. (2017). Effects of air pollution on the skin: A review. *Indian Journal of Dermatology, Venereology and Leprology*, 83(5), 415–423. <https://doi.org/10.4103/0378-6323.200328>
- [26] Invernizzi, G., Ruprecht, A., Mazza, R., et al. (2004). Particulate matter from tobacco versus diesel car exhaust: An educational perspective. *Tobacco Control*, 13(3), 219–221. <https://doi.org/10.1136/tc.2004.009885>
- [27] Blois, M. S. (1958). Antioxidant determinations by the use of a stable free radical. *Nature*, 181(4617), 1199–1200. <https://doi.org/10.1038/1811199a0>
- [28] Regazzetti, C., Sormani, L., Debayle, D., et al. (2018). Melanocytes sense blue light and regulate pigmentation through opsin-3. *Journal of Investigative Dermatology*, 138(1), 171–178. <https://doi.org/10.1016/j.jid.2017.08.044>
- [29] Ascenso, A., Ribeiro, H. M., Marques, H. C., & Simões, S. (2011). Topical delivery of antioxidants. *Current Drug Delivery*, 8(6), 640–660. <https://doi.org/10.2174/156720111796181161>
- [30] Kaur, I. P., Kapila, M., & Agrawal, R. (2007). Role of novel delivery systems in developing topical antioxidants as therapeutics to combat photoaging. *Ageing Research Reviews*, 6(4), 271–288. <https://doi.org/10.1016/j.arr.2007.03.003>
- [31] Burke, K. E. (2018). Mechanisms of aging and development: A new understanding of environmental damage to the skin and prevention with topical antioxidants. *Mechanisms of Ageing and Development*, 172, 123–130. <https://doi.org/10.1016/j.mad.2017.10.007>
- [32] Duarah, S., Durai, R. D., & Narayanan, V. B. (2017). Nanoparticle-in-gel system for delivery of vitamin C for topical application. *Drug Delivery and Translational Research*, 7(6), 750–760. <https://doi.org/10.1007/s13346-017-0445-x>
- [33] Krutmann, J., Liu, W., Li, L., et al. (2014). Pollution and skin: From epidemiological and mechanistic studies to clinical implications. *Journal of Dermatological Science*, 76(3), 163–168. <https://doi.org/10.1016/j.jdermsci.2014.02.00>
- [34] Jin, S. P., Li, Z., Choi, E. K., et al. (2018). Urban particulate matter in air

- pollution penetrates into the barrier-disrupted skin and produces ROS-dependent cutaneous inflammatory response in vivo. *Journal of Dermatological Science*, 91(2), 175–183. <https://doi.org/10.1016/j.jdermsci.2018.01.007>
- [35] Mancebo, S. E., & Wang, S. Q. (2015). Recognizing the impact of ambient air pollution on skin health. *Journal of the European Academy of Dermatology and Venereology*, 29(12), 2326–2332. <https://doi.org/10.1111/jdv.1308>
- [36] Soeur, J., Belaïdi, J. P., Chollet, C., et al. (2017). Photo-pollution stress in skin: Traces of pollutants (PAH and particulate matter) impair redox homeostasis in keratinocytes exposed to UVA1. *Journal of Dermatological Science*, 86(2), 162–169. <https://doi.org/10.1016/j.jdermsci.2017.08.003>
- [37] McDaniel, D., Farris, P., & Valacchi, G. (2018). Atmospheric skin aging – Contributors and inhibitors. *Journal of Cosmetic Dermatology*, 17(2), 124–137. <https://doi.org/10.1111/jocd.12460>
- [38] Latha, M. S., Martis, J., Shobha, V., et al. (2013). Sunscreening agents: A review. *Journal of Clinical and Aesthetic Dermatology*, 6(8), 16–26.
- [39] Heurung, A. R., Raju, S. I., & Warshaw, E. M. (2014). Adverse reactions to sunscreen agents: Epidemiology, responsible irritants and allergens, clinical characteristics, and management. *Dermatitis*, 25(5), 289–326. <https://doi.org/10.1097/DER.0000000000000056>
- [40] Ruszkiewicz, J. A., Pinkas, A., Ferrer, B., Peres, T. V., Tsatsakis, A., & Aschner, M. (2017). Neurotoxic effect of active ingredients in sunscreen products: A contemporary review. *Toxicology Reports*, 4, 245–259. <https://doi.org/10.1016/j.toxrep.2017.04.003>
- [41] Drakaki, E., Dessinioti, C., & Antoniou, C. (2014). Air pollution and the skin. *Frontiers in Environmental Science*, 2, 11. <https://doi.org/10.3389/fenvs.2014.0001>
- [42] Nguyen, T., Nioi, P., & Pickett, C. B. (2009). The Nrf2-antioxidant response element signaling pathway and its activation by oxidative stress. *Journal of Biological Chemistry*, 284(20), 13291–13295. <https://doi.org/10.1074/jbc.R900010200>
- [43] Ma, Q., & Lu, A. Y. (2007). CYP1A induction and human risk assessment: An evolving tale of in vitro and in vivo studies. *Drug Metabolism and Disposition*, 35, 1009–1016. <https://doi.org/10.1124/dmd.107.01582>
- [44] Catalá, M., Gasulla, F., Pradas Del Real, A. E., García-Breijo, F., Reig-Armiñana, J., & Barreno, E. (2013).

The organic air pollutant cumene hydroperoxide interferes with NO antioxidant role in rehydrating lichen. *Environmental Pollution*, 179, 277–284. <https://doi.org/10.1016/j.envpol.2013.04.015>

[45] Wang, S., Virmani, P., & Lim, H. (2016). Consumer acceptability and compliance: The next frontier in sunscreen innovation. *Photodermatology, Photoimmunology & Photomedicine*, 32(1), 55–56. <https://doi.org/10.1111/phpp.12211>

[46] Prescott, S. L., Larcombe, D. L., Logan, A. C., West, C., Burks, W., Caraballo, L., Levin, M., Etten, E. V., Horwitz, P., Kozyrskyj, A., et al. (2017). The skin microbiome: Impact of modern environments on skin ecology, barrier integrity, and systemic immune programming. *World Allergy Organization Journal*, 10, 29. <https://doi.org/10.1186/s40413-017-0199-9>

[47] Iwase, T., Uehara, Y., Shinji, H., Tajima, A., Seo, H., Takada, K., Agata, T., & Mizunoe, Y. (2010). *Staphylococcus epidermidis* Esp inhibits *Staphylococcus aureus* biofilm formation and nasal colonization. *Nature*, 465, 346–349. <https://doi.org/10.1038/nature09074>

[48] Zipperer, A., Konnerth, M. C., Laux, C., Berscheid, A., Janek, D., Weidenmaier, C., Burian, M.,

Schilling, N. A., Slavetinsky, C., Marschal, M., et al. (2016). Human commensals producing a novel antibiotic impair pathogen colonization. *Nature*, 535, 511–516. <https://doi.org/10.1038/nature18623>

[49] Nakatsuji, T., Chen, T. H., Narala, S., Chun, K. A., Two, A. M., Yun, T., Shafiq, F., Kotol, P. F., Bouslimani, A., Melnik, A. V., et al. (2017). Antimicrobials from human skin commensal bacteria protect against *Staphylococcus aureus* and are deficient in atopic dermatitis. *Science Translational Medicine*, 9, 378. <https://doi.org/10.1126/scitranslmed.ah4680>

[50] Zheng, Y., Hunt, R. L., Villaruz, A. E., Fisher, E. L., Liu, R., Liu, Q., Cheung, G. Y. C., Li, M., & Otto, M. (2022). Commensal *Staphylococcus epidermidis* contributes to skin barrier homeostasis by generating protective ceramides. *Cell Host & Microbe*, 30, 301–313. <https://doi.org/10.1016/j.chom.2022.01.009>

[51] Cau, L., Williams, M. R., Butcher, A. M., Nakatsuji, T., Kavanaugh, J. S., Cheng, J. Y., Shafiq, F., Higbee, K., Hata, T. R., Horswill, A. R., et al. (2021). *Staphylococcus epidermidis* protease EcpA can be a deleterious component of the skin microbiome in atopic dermatitis. *The Journal of Allergy and Clinical Immunology*,

147, 955–966. <https://doi.org/10.1016/j.jaci.2020.08.016>

- [52] Lee, H. J., Jeong, S. E., Lee, S., Kim, S., Han, H., & Jeon, C. O. (2018). Effects of cosmetics on the skin microbiome of facial cheeks with different hydration levels. *MicrobiologyOpen*, 7, e00557. <https://doi.org/10.1002/mbo3.557>

- [53] Ghosal, D., Ghosh, S., Dutta, T. K., & Ahn, Y. (2016). Current state of knowledge in microbial degradation of polycyclic aromatic hydrocarbons (PAHs): A review. *Frontiers in Microbiology*, 7, 1369. <https://doi.org/10.3389/fmicb.2016.01369>

